SYNTHETIC COMMUNICATIONS[®] Vol. 34, No. 19, pp. 3449–3453, 2004

Conversion of α-Amino Acids into Nitriles by Oxidative Decarboxylation with Trichloroisocyanuric Acid

Gene A. Hiegel,* Justin C. Lewis, and Jason W. Bae

Department of Chemistry and Biochemistry, California State University, Fullerton, California, USA

ABSTRACT

Trichloroisocyanuric acid oxidation of α -amino acids in water or methanol in the presence of pyridine produces nitriles with one less carbon in good yields and of high purity.

Key Words: α-Amino acids; Nitriles; Oxidative decarboxylation; Oxidation; Decarboxylation; Preparation; Synthesis; Trichloroisocyanuric acid.

Amino acid oxidations have been carried out with a variety of oxidizing agents to produce a variety of products. Silver oxide oxidation produces car-

3449

DOI: 10.1081/SCC-200030958 Copyright © 2004 by Marcel Dekker, Inc. 0039-7911 (Print); 1532-2432 (Online) www.dekker.com

Request Permissions / Order Reprints

^{*}Correspondence: Gene A. Hiegel, Department of Chemistry and Biochemistry, California State University, Fullerton, CA 92834, USA; E-mail: ghiegel@fullerton. edu.

boxylic acids with one less carbon.^[1] Electrochemical oxidation in the presence of triphenylphosphine gives α -amino aldehydes.^[2] Silver catalyzed oxidation by persulfate generates aldehydes with one less carbon.^[3] The reaction with chlorine produces a mixture of aldehydes and nitriles with one less carbon.^[4] A search of the literature did not reveal any synthetic procedures for the conversion of α -amino acids effectively into nitriles. We wish to report that α -amino acids are converted cleanly and efficiently into nitriles by means of an oxidative decarboxylation reaction carried out with trichloroisocyanuric acid [TCICA; 1,3,5-trichloro-1,3,5-triazine-2,4,6-(1H,3H,5H)-trione; C₃Cl₃N₃O₃] in water or methanol in the presence of pyridine Eq. (1). The reaction is exothermic, and therefore, TCICA is added slowly while using a dry-ice condenser for the lower molecular weight products. Yields of the more volatile nitriles were considerably lower without the dry-ice condenser. The results are shown in Table 1.



L-Isoleucine [(2S,3S)-(+)-2-amino-3-methylpentanoic acid] was converted into (S)-(+)-2-methylbutyronitrile with a specific rotation of +37.46° for the purest fraction. This nitrile has also been prepared from the corresponding aldehyde, from the oxime of the aldehyde, and from the corresponding carboxylic acid.^[5] The highest observed specific rotation was +32.7°, although the estimated maximum specific rotation was 36.3°. Oxidation of isoleucine with TCICA appears to be the best method for the synthesis of (S)-(+)-2-methylbutyronitrile considering the simplicity of the reaction, the availability of isoleucine, and the high optical purity obtained.

EXPERIMENTAL

All reagents were used as received. Valine, leucine, isoleucine, norleucine, phenylalanine, phenylglycine, glutaric acid, isobutyronitrile, isovaleronitrile, 2-methylbutyronitrile, valeronitrile, benzonitrile, and methanol were obtained from Aldrich. Pyridine and phenylacetonitrile were obtained from J. T. Baker. The TCICA (99%) was obtained from Chem Lab Products. All products except for methyl 3-cyanopropanoate were compared with authentic samples by means of ¹H FT-NMR and Fourier-transform infrared (FT-IR) spectra and GC retention time. 3-Cyanopropanoic acid was converted into methyl 3-cyanopropanoate before isolation and characterized by means of the IR spectrum and boiling point.

Amino acid	Nitrile	Yield	Boiling point	GC purity
ОН	C≡N	58.4% ^a	102.7–103.0°C	99.1%
о ОН	C≡N	79.3% ^a	128.0–128.5°C	97.9%
O OH	C=N	88.0% ^a	124.0-125.0°C	97.4%
№H₂ ОН	C=N	84.2% ^a	139.5–140.0°C	98.9%
NH2 О ОН	C≡N	66.1% ^a	190.0°C	99.3%
NH ₂ O O O H	C≡N	77.0% ^b	c	99.9%
HO OH	CH ₃ O C≡N	61.4% ^b	104.0–105.2°C (13.4–14.2 torr) ^d	96.5%

Table 1. Conversion of α -amino acids into nitriles.

^aReaction run in water.

^bReaction run in methanol.

^cPurified by flash chromatography.

^dConverted to the methyl ester before isolation.

¹H FT-NMR spectra were recorded in CDCl₃ using an Anasazi-modified Varian EFT 90 MHz spectrometer. The FT-IR spectra were recorded using a Perkin Elmer 1650 spectrometer. GC analyses were carried out with a Hewlett-Packard 5890 Series II on a 6 ft \times 1/8 in 10% Carbowax 20-M column. Optical rotations were recorded using a Rudolph Instruments Model 51 polarimeter equipped with a sodium vapor lamp.

Conversion of L-Isoleucine into (S)-(+)-2-Methylbutyronitrile

L-isoleucine, 15.300 g (0.1166 mol), 120 mL water, 29.05 g (0.3673 mol) pyridine, and a magnet were placed in a 300-mL three-neck round bottom

flask fitted with a dry-ice condenser, a stopper, and a solid addition funnel containing 23.362 g (0.1000 mol, 0.3000 eq) TCICA. While stirring, TCICA was added over a period of 30 min. The reaction was exothermic. After 3 hr additional stirring at room temperature, solid NaHSO₃ was added in order to destroy excess TCICA until a negative test with iodide-starch paper was obtained. After filtration to remove the cyanuric acid that formed and washing the solid with ether, the mixture was acidified with HCl and the aqueous phase was saturated with NaCl. The mixture was extracted with ether (4 × 50 mL), dried over MgSO₄, and filtered. The ether was removed by distillation through a concentric tube column and the amine distilled to give several fractions of (S)-(+)-2-methylbutyronitrile, bp 124.0–125.0°C. Fractions 3–6, 8.532 g (88.0%), had a purity of 97.4% by GC. Fractions 4–6, 7.618 g (78.6%) had a purity of 99.2% by GC. Fraction 5, 4.904 g (50.6%), had a purity of 99.5% by GC: $[\alpha]_D^{20} + 37.46^\circ$ (neat). The FT-IR, ¹H FT-NMR, and GC retention time were identical to an authentic sample.

Conversion of L-Glutaric Acid into Methyl 3-Cyanopropanoic Acid

Glutaric acid, 15.655 g (0.1064 mol), 75 mL methanol, 17.67 g. (0.2233 mol) pyridine, and a magnet were placed in a 300-mL three-neck round bottom flask fitted with a dry-ice condenser, a stopper, and a solid addition funnel containing 21.098 g (0.09077 mol, 0.2723 eq) TCICA. While stirring, TCICA was added over a period of 45 min. The reaction was exothermic. After 1 hr 50 min additional stirring at room temperature, solid NaHSO₃ was added in order to destroy excess TCICA until a negative test with iodidestarch paper was obtained. After filtration to remove the cyanuric acid that formed and washing the solid with methanol, 7 mL conc H₂SO₄ was added to the filtrate and the solution was refluxed for 3 hr in order to form the methyl ester. The methanol was removed using a rotary evaporator and 25 mL of water was added, and the mixture was extracted with ether $(3 \times 100 \text{ mL})$. The ether solution was dried over MgSO₄, and after filtration, the ether was removed using a rotary evaporator and the residue distilled using a concentric tube column to give 7.338 g. (61.4%) [bp 104.0-105.2°C (13.4-14.2 torr)] methyl 3-cyanopropanoate with a purity by GC of 96.5% [reported bp 95–97°C (8 torr)];^[6] FT-IR (film) 2252 (CN), 1739 (CO), 1293, 1254, 1209, 1179, 1058, 1023, 1001, 981, 946, 878, 838, 784, 705 cm⁻¹ [reported (film) 2246, 1735, 1286, 1247, 1202, 1172, 1052, 1017, 996, 975, 940, 871, 833, 778, 697 cm^{-1} ;^[7] ¹H FT-NMR (CDCl₃) δ 3.76 (s, 3, OCH₃), 2.70 (s, 4, CH₂CH₂).

ACKNOWLEDGMENT

This research was supported in part by a grant from the CSU Special Fund for Research, Scholarship, and Creative Activity to G. A. H. and funds from the Southern California Academy of Sciences Research Training Program to J. W. B.

REFERENCES

- Clarke, T.G.; Hampson, N.A.; Lee, J.B.; Morley, J.R.; Scanlon, B. Argentic oxide oxidations of organic compounds. Tetrahedron Lett. 1968, (54), 5685–5688.
- 2. Maeda, H.; Maki, T.; Ohmori, H. One-step transformation of α -amino acids to α -amino aldehydes effected by electrochemical oxidation of Ph₃P. Tetrahedron Lett. **1992**, *33* (10), 1347–1350.
- 3. Bacon, R.G.R.; Hanna, W.J.W.; Stewart, D. Oxidation by persulfate. Part V. Silver-catalyzed oxidation of secondary aliphatic amines and α -amino acids. J. Chem. Soc. (C) **1966**, 1388–1389.
- Ishikawa, S.; Fujino, H.; Yasuda, K.; Shigemori, N. Products at the reaction of amino acids in aqueous solution with chlorine. Suishitsu Odaku Kenkyu 1986, 9 (12), 786–792.
- Botteghi, C.; Chelucci, G.; Marchetti, M. Synthesis of optically active aliphatic nitriles from aldehydes and acids. Synth. Commun. 1982, *12* (1), 25–33.
- 6. McElvain, S.M.; Schroeder, J.P. Orthoesters and related compounds from malono- and succinonitriles. J. Am. Chem. Soc. **1949**, *71*, 40–46.
- Simons, W.W. Ed. *The Sadtler Handbook of Infrared Spectra*; Sadtler Research Laboratories, Inc.: Philadelphia, PA, 1978; spectrum no. 2737.

Received in the USA May 27, 2004

Copyright of Synthetic Communications is the property of Marcel Dekker Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.